

# Insights into calcium-sensing receptor trafficking and biased signalling by studies of calcium homeostasis

Gorvin, Caroline

DOI:

[10.1530/JME-18-0049](https://doi.org/10.1530/JME-18-0049)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Gorvin, C 2018, 'Insights into calcium-sensing receptor trafficking and biased signalling by studies of calcium homeostasis', *Journal of Molecular Endocrinology*. <https://doi.org/10.1530/JME-18-0049>

[Link to publication on Research at Birmingham portal](#)

## Publisher Rights Statement:

Disclaimer: this is not the definitive version of record of this article. This manuscript has been accepted for publication in Journal of Molecular Endocrinology, but the version presented here has not yet been copy-edited, formatted or proofed. Consequently, Bioscientifica accepts no responsibility for any errors or omissions it may contain. The definitive version is now freely available at: <http://dx.doi.org/10.1530/JME-18-0049>, pub 2018

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1    **Insights into calcium-sensing receptor trafficking and biased signalling by studies of calcium**  
2    **homeostasis**

3    Caroline M. Gorvin<sup>a,b</sup>

4

5    **Author affiliations**

6    <sup>a</sup>Institute of Metabolism and Systems Research (IMSR), University of Birmingham, Birmingham,  
7    B15 2TT.

8    <sup>b</sup>Centre for Endocrinology, Diabetes and Metabolism (CEDAM), Birmingham Health Partners,  
9    Birmingham, B15 2TH, UK

10    Email: [C.Gorvin@bham.ac.uk](mailto:C.Gorvin@bham.ac.uk)

11    Telephone: +44 (0)121 414 6857

12

13    **Word Count:** words (4000-8000 words (excluding references and Figure Legends))

14    **Short Title:** CaSR trafficking and signalling

15    **Keywords:** calcium-sensing receptor, adaptor protein-2, familial hypocalciuric hypercalcaemia,  
16    autosomal dominant hypocalcaemia, biased signalling

17

18

19

20

21

22

## 23 ABSTRACT

24 The calcium-sensing receptor (CaSR) is a class C G-protein coupled receptor (GPCR) that detects  
25 extracellular calcium concentrations, and modulates parathyroid hormone secretion and urinary  
26 calcium excretion to maintain calcium homeostasis. The CaSR utilises multiple heterotrimeric G-  
27 proteins to mediate signalling effects including: activation of intracellular calcium release; mitogen-  
28 activated protein kinase (MAPK) pathways; membrane ruffling; and inhibition of cAMP production.  
29 By studying germline mutations in the CaSR and proteins within its signalling pathway that cause  
30 hyper- and hypocalcaemic disorders, novel mechanisms governing GPCR signalling and trafficking  
31 have been elucidated. This review focusses on two recently described pathways that provide novel  
32 insights into CaSR signalling and trafficking mechanisms. The first, identified by studying a CaSR  
33 gain-of-function mutation that causes autosomal dominant hypocalcaemia (ADH), demonstrated a  
34 structural motif located between the third transmembrane domain and the second extracellular loop of  
35 the CaSR that mediates biased signalling by activating a novel  $\beta$ -arrestin-mediated G-protein-  
36 independent pathway. The second, in which the mechanism by which adaptor protein-2  $\sigma$ -subunit  
37 (AP2 $\sigma$ ) mutations cause familial hypocalciuric hypercalcaemia (FHH) was investigated, demonstrated  
38 that AP2 $\sigma$  mutations impair CaSR internalisation and reduce multiple CaSR-mediated signalling  
39 pathways. Furthermore, these studies showed that the CaSR can signal from the cell surface using  
40 multiple G-protein pathways, whilst sustained signalling is mediated only by the  $G_{q/11}$  pathway. Thus,  
41 studies of FHH and ADH associated mutations have revealed novel steps by which CaSR mediates  
42 signalling and compartmental bias, and these pathways could provide new targets for therapies for  
43 patients with calcaemic disorders.

44     **1       Introduction to the CaSR**

45     Extracellular calcium ( $\text{Ca}^{2+}_e$ ) is required for diverse biological functions ranging from blood  
 46     coagulation, mineralisation of bone matrix, muscle contraction, and hormone secretion (Brown 1991).  
 47     Thus, calcium concentrations within the blood are tightly regulated. The parathyroid gland plays an  
 48     essential role in calcium homeostasis by detecting  $\text{Ca}^{2+}_e$  in the blood, and in response to  
 49     hypocalcaemia, secretes the parathyroid hormone (PTH) to normalize serum calcium concentrations.  
 50     PTH achieves this by: enhancing bone resorption; activating calcium reabsorption at the kidneys; and  
 51     stimulating the synthesis of 1,25-dihydroxyvitamin  $\text{D}_3$ , which mobilises intestinal calcium absorption  
 52     (Fig. 1A) (Riccardi and Brown 2010). The net effect of these three pathways is to increase  $\text{Ca}^{2+}_e$ ,  
 53     which provides feedback inhibition to the parathyroid gland, to suppress PTH secretion (Conigrave  
 54     and Ward 2013). The parathyroid is able to detect  $\text{Ca}^{2+}_e$  concentrations in the blood using the cell-  
 55     surface expressed calcium-sensing receptor (CaSR), a class C G-protein coupled receptor (GPCR), for  
 56     which  $\text{Ca}^{2+}$  is the major ligand (Conigrave and Ward 2013; Riccardi and Brown 2010).

57     The CaSR is a 1078 amino acid protein that exists at cell surfaces as a disulphide-linked homodimer  
 58     (Ward, et al. 1998), although it is capable of forming heterodimers (Chang, et al. 2007; Gama, et al.  
 59     2001). The CaSR has a large extracellular domain, which was recently crystallized by two  
 60     independent labs, and consists of a bilobed venus fly-trap ligand binding domain (VFTD), and a  
 61     cysteine-rich domain (CRD) (Geng, et al. 2016; Zhang, et al. 2016) (Fig. 1B).  $\text{Ca}^{2+}_e$  binds between the  
 62     two lobes (lobe 1 and lobe 2) of the VFTD, which initiates a conformational change, facilitating lobe  
 63     2-lobe 2 dimerisation and allowing the cysteine-rich domains to interact (Geng et al. 2016; Zhang et  
 64     al. 2016). These conformational changes are predicted to re-orientate the seven transmembrane (TM)  
 65     domain (Geng et al. 2016), and consequently activate the associated G-proteins and initiate signal  
 66     transduction (Standfuss, et al. 2011). The CaSR has been reported to couple to multiple G-protein  
 67     subtypes, but predominantly signals by: the  $\text{G}_{i/o}$  pathway, to suppress cAMP and activate mitogen-  
 68     activated protein kinase (MAPK) cascades (Kifor, et al. 2001; Thomsen, et al. 2012); the  $\text{G}_{q/11}$ -  
 69     phospholipase C (PLC)-mediated pathway, to generate inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ) and  
 70     diacylglycerol, that activate intracellular calcium ( $\text{Ca}^{2+}_i$ ) mobilisations (Hofer and Brown 2003) and

MAPK pathways, respectively. The CaSR can also couple to a G-protein-independent mechanism involving  $\beta$ -arrestin proteins to also activate MAPK signalling (Thomsen et al. 2012) (Fig. 2).

73

## 2 Human disorders of the CaSR provide insights into receptor signalling mechanisms

The importance of the CaSR in the regulation of  $\text{Ca}^{2+}_e$  is highlighted in patients with germline mutations of the receptor that have disorders of calcium homeostasis. Heterozygous loss-of-function mutations in the CaSR lead to the autosomal dominant condition familial hypocalciuric hypercalcaemia (FHH) (Pollak, et al. 1993), which is characterised by lifelong mild-to-moderate hypercalcaemia, normal or mildly raised serum PTH levels and low urinary calcium excretion (Firek, et al. 1991; Marx 2015). FHH is considered to be a benign disorder as most patients are asymptomatic, but it can be associated with chondrocalcinosis and pancreatitis in some cases (Hannan and Thakker 2013; Pearce, et al. 1996a; Volpe, et al. 2009). Homozygous and compound heterozygous loss-of-function mutations of the CaSR are the common cause of neonatal severe hyperparathyroidism (NSHPT), which is characterised by marked elevations in serum calcium and PTH, failure to thrive and hyperparathyroid bone disease (Chattopadhyay and Brown 2006; Hannan and Thakker 2013; Pollak et al. 1993). On occasion, heterozygous CaSR mutations may lead to an NSHPT-like phenotype, and this is largely due to the dominant-negative nature of the mutant receptor on the wild-type CaSR in these cases (Obermannova, et al. 2009; Pearce, et al. 1995).

Gain-of-function mutations of the CaSR cause autosomal dominant hypocalcaemia (ADH) characterised by mild-to-moderate hypocalcaemia and inappropriately low or normal PTH concentrations (Hannan and Thakker 2013; Pearce, et al. 1996b). Up to 50% of patients present with hypocalcaemic symptoms of paraesthesia, carpopedal spasms, seizures, and ectopic calcification of the kidneys and basal ganglia (Hannan and Thakker 2013; Pearce et al. 1996b). Some patients with gain-of-function mutations in CaSR may have Bartter syndrome type 5, which is characterised by renal salt wasting, hypokalaemia, hyperreninaemia and hyperaldosteronaemia (Vargas-Poussou, et al. 2002; Watanabe, et al. 2002).

## 97    **2.1    Biased signalling of the CaSR**

98    Functional studies in HEK293 cells have demonstrated that disease-causing mutations may influence  
 99    CaSR signalling responses in a biased manner (Leach, et al. 2012). Despite the capability of the CaSR  
 100    to signal via multiple signalling pathways, *in vitro* studies in HEK293 cells have shown that it  
 101    preferentially couples to the  $\text{Ca}^{2+}_i$  signalling pathway (Leach et al. 2012). In contrast, disease-causing  
 102    CaSR mutants have been shown to switch this preferential signalling, with some FHH1-causing  
 103    mutations signalling equally via the  $\text{Ca}^{2+}_i$  and MAPK pathways, or predominantly via MAPK  
 104    pathways; while many ADH1-associated mutants couple more strongly to  $\text{Ca}^{2+}_i$  (Leach et al. 2012).  
 105    Furthermore, studies of positive and negative allosteric CaSR modulating compounds, revealed they  
 106    too can mediate a biased signalling response, with both classes of drugs influencing  $\text{Ca}^{2+}_i$  to a greater  
 107    extent than ERK1/2 phosphorylation (Leach, et al. 2013). Thus, these findings established that  
 108    agonist-induced CaSR signalling may occur in a biased manner, although the GPCR structural motifs  
 109    mediating ligand-dependent bias were not described.

110

## 111    **2.2    Novel insights into mechanisms by which CaSR can mediate signalling bias**

112    Recently, a novel ADH1-associated CaSR mutation, Arg680Gly, has provided some insights into the  
 113    molecular mechanisms mediating signalling bias (Gorvin, et al. 2018a). *In vitro* analyses showed that  
 114     $\text{Ca}^{2+}_i$  signalling in cells expressing the Arg680Gly mutation was not different to that observed in wild-  
 115    type cells, in contrast to previously reported ADH1-associated CaSR mutants (Gorvin et al. 2018a).  
 116    However, the Arg680Gly mutation did enhance MAPK signalling (Gorvin et al. 2018a). Furthermore,  
 117    this type of signalling bias, in which an ADH1 mutation enhances MAPK signalling, but does not  
 118    affect  $\text{Ca}^{2+}_i$  has not previously been described and provided an opportunity to explore the role of the  
 119    Arg680 residue in CaSR structure-function. As MAPK acts as a convergence pathway for multiple  
 120    CaSR signalling pathways (Fig. 2) each was investigated using a single reporter assay measuring  
 121    luciferase upstream of a serum-response element (SRE), which acts as a measure for MAPK-mediated  
 122    transcription (Gorvin, et al. 2017b). By applying compounds that specifically block either the  $\text{G}_{q/11}$

((YM-245890 and UBO-QIC), the  $G_{i/o}$  (pertussis toxin) or the  $\beta$ -arrestin ( $\beta$ -arrestin-1 and  $\beta$ -arrestin-2 targeting siRNA) pathways, it was shown that the Arg680Gly mutation enhanced MAPK signalling by a  $\beta$ -arrestin1/2-mediated pathway (Gorvin et al. 2018a; Schrage, et al. 2015; Takasaki, et al. 2004).

The structural location of the mutant residue within the CaSR transmembrane domain (TMD) region provides some insights into the likely mechanism mediating this bias. Homology modelling of the CaSR TMD, based upon the structure of the closely related human metabotropic glutamate receptor 1 (mGluR1) (Gorvin et al. 2018a; Hu, et al. 2005), predicts that the Arg680 residue lies at the extracellular side of TM3, and that the residue forms salt bridge connections with the side chain of adjacent residues in extracellular loop 2 (ECL2) (Glu767) or TM7 (Glu837) (Fig. 3). Such connections between residues within TMDs or with ECLs of GPCRs are known to be important in receptor activity. For example, TM3 and TM6 of the  $\beta$ 2AR forms ionic interactions which lock the receptor in an inactive state, and conformational changes within these helices governs G-protein coupling at the cytoplasmic face of the receptor (Ballesteros, et al. 2001; Rasmussen, et al. 2007). Furthermore, disruption of a salt bridge in  $\beta$ 2-AR allows lateral displacement of TM3 away from TM4 and TM5, facilitating  $\beta$ -arrestin binding (Shukla, et al. 2014), and it was therefore hypothesised that the CaSR Arg680Gly mutation may break a salt-bridge between TM3 and ECL2 or TM7 allowing  $\beta$ -arrestin to bind more readily at the CaSR cytoplasmic face (Gorvin et al. 2018a). The importance of the Arg680, Glu767 and Glu837 residues in the CaSR has previously been recognised. Mutation of Glu767 and Glu837 has been shown to increase signaling by the CaSR (Hu, et al. 2006; Hu et al. 2005; Hu, et al. 2002; Uckun-Kitapci, et al. 2005), and previous structural homology models that investigated the binding of allosteric modulator drugs with the CaSR demonstrated that the three residues form critical contacts for drug binding (Miedlich, et al. 2004; Petrel, et al. 2004), indicating their importance in CaSR structure-function. To investigate the hypothesis that a salt-bridge is disrupted by mutation of Arg680, a series of mutations were engineered at the Arg680, Glu767 and Glu837 residues. First, Glu767 and Glu837 were mutated to Arg residues, to introduce unfavourable electrostatic interactions. Under these conditions, the Glu767Arg engineered mutant was shown to

enhance MAPK activity in a similar way to the Arg680Gly ADH1 mutant, a response that was also  $\beta$ -arrestin sensitive (Gorvin et al. 2018a). In contrast, responses in cells expressing the CaSR Glu837Arg engineered mutant were similar to those in cells expressing WT CaSR (Gorvin et al. 2018a). To confirm that a salt-bridge is required between residues 680 and 767 in the CaSR, a double mutant was generated by mutating the Glu767Arg mutant receptor with an additional missense mutation of Arg680 to Glu680. This should allow the salt-bridge between the two residues to reform. Indeed, in this double mutant (Glu680-Arg767) CaSR, MAPK signalling was restored to levels similar to those seen in wild-type cells (Gorvin et al. 2018a).

158

Thus, these studies have revealed some of the important structural motifs of CaSR that mediate signalling bias, and discovery of this novel  $\beta$ -arrestin-specific pathway may help facilitate the development of targeted therapeutics for CaSR. Furthermore, it demonstrates the importance of investigating multiple signalling outputs downstream of CaSR, to ensure that potentially disease-causing mutations are not classified as benign polymorphisms (Gorvin et al. 2018a).

164

### 3      **Insights from human genetic mutations in components of the CaSR signalling pathway**

Both FHH and ADH are genetically heterozygous conditions with mutations in the CaSR accounting for approximately 65% and 70% of cases, respectively (Hannan, et al. 2012; Nesbit, et al. 2013a). Two further genetically distinct forms of FHH, and one further distinct form of ADH, have been described (Mannstadt, et al. 2013; Nesbit et al. 2013a; Nesbit, et al. 2013b). Heterozygous loss- and gain-of-function germline mutations in the  $\alpha$ -subunit of the G-protein 11 ( $G\alpha_{11}$ ), a component of the CaSR signalling pathway, give rise to FHH type-2 (FHH2) and ADH type-2 (ADH2), respectively (Mannstadt et al. 2013; Nesbit et al. 2013a). Only four FHH2 and six ADH2 mutations have been described to date, and therefore these mutations account for only a small number of cases of these disorders (Gorvin, et al. 2016; Gorvin, et al. 2017a; Li, et al. 2014; Mannstadt et al. 2013; Nesbit et al. 2013a; Piret, et al. 2016). FHH type-3 (FHH3) is due to mutations in the sigma subunit of the adaptor



protein-2 (AP2 $\sigma$ ), which plays a fundamental role in clathrin-mediated endocytosis of transmembrane proteins, such as GPCRs (Nesbit et al. 2013b). The AP2 $\sigma$  protein is ubiquitously expressed and clathrin-mediated endocytosis is a critical cellular process; however, FHH is a largely benign condition, and the phenotypes observed in FHH3 patients are largely CaSR-specific. Thus, by studying the AP2 $\sigma$  mutations identified in FHH3, which have been described in a single residue (Hannan, et al. 2015; Nesbit et al. 2013b), novel insights into the trafficking and signalling mechanisms of CaSR have been elucidated, and indicate an important interplay between these two processes (Gorvin, et al. 2018b).

### 3.1 Trafficking of the CaSR

The cell surface expression of the CaSR is important for detecting extracellular ligand and signalling by the receptor, and therefore the plasma membrane expression of CaSR is carefully regulated. This involves multiple pathways including: receptor synthesis and secretion, trafficking to the plasma membrane, and removal of the CaSR from the cell surface by endocytosis.

#### 3.1.1 Regulation of cell surface expression by functional desensitisation

In contrast to most GPCRs, which undergo agonist-dependent desensitisation by phosphorylation and/or by  $\beta$ -arrestin proteins, functional desensitisation appears to have only a minimal impact on CaSR expression, and there are inconsistencies between findings in these studies (Bouschet, et al. 2005; Lorenz, et al. 2007; Pi, et al. 2005; Thomsen et al. 2012). Initial studies of the CaSR showed that several proteins desensitise the receptor. These include: PKC, which phosphorylates the receptor following activation of signalling pathways and recruits  $\beta$ -arrestin; G-protein regulatory kinase (GRK) 2, which binds to  $G\alpha_q$  and inhibits its signalling; and, GRK4 which phosphorylates CaSR to facilitate desensitisation (Lorenz et al. 2007; Pi et al. 2005). Furthermore, GRKs and  $\beta$ -arrestin have independent functions as combined treatment of cells with these proteins enhances desensitisation (Lorenz et al. 2007). The traditional view of GPCR desensitisation and internalisation was one in

which the receptor is phosphorylated and  $\beta$ -arrestin recruited, followed by internalisation of the receptor by clathrin-mediated endocytosis, which is facilitated by interactions between  $\beta$ -arrestin and AP2 (Shukla, et al. 2011). While studies of CaSR showed that  $\beta$ -arrestin can be recruited, this did not enhance receptor internalisation, which only occurred when cells were treated with high concentrations of  $\text{Ca}^{2+}_e$  (10mM) (Lorenz et al. 2007).  $\beta$ -arrestin is also now recognised to have another function, as a scaffold protein that facilitates signalling either at the plasma membrane, or for some GPCRs, at the endosome (Shukla et al. 2014). This function of  $\beta$ -arrestin has been recognised for CaSR in some studies, providing seemingly contradictory information to that in studies of functional desensitisation. Thus, treatment of cells with dominant-negative forms of  $\beta$ -arrestin1 or  $\beta$ -arrestin2, or with siRNA targeting  $\beta$ -arrestin1 or  $\beta$ -arrestin2, reduces the pERK and membrane ruffling signals downstream of CaSR (Bouschet et al. 2005; Gorvin et al. 2018b; Thomsen et al. 2012). Further studies are required to determine whether the discrepancies within these data sets are due to experimental differences, differences in cell type, or if both desensitisation and enhanced signalling occur downstream of CaSR, but at different spatial or temporal points.

### **3.1.2 Regulation of cell surface expression by the secretory pathway and agonist-driven insertional signalling (ADIS)**

The cell surface expression of CaSR, and mechanisms by which the receptor is synthesised and exported to the plasma membrane, are generally well understood. The CaSR is synthesised at the ER where it forms homodimers (Pidashveva, et al. 2006) and undergoes quality control and immature glycosylation steps prior to progression to the Golgi (Fan, et al. 1997). At the Golgi, the CaSR undergoes further maturation. This forward trafficking through the secretory pathway involves a number of regulatory proteins that have been well described in previous reviews (Breitwieser 2013; Huang, et al. 2011). The core-glycosylated CaSR is then retained within pre-plasma membrane compartments (Fan et al. 1997). Therefore CaSR is retained intracellularly within two large intracellular reserves: one at the ER and one at pre-plasma membrane compartments (Breitwieser 2012).

229

230 The large pool of fully mature CaSR plays a unique role in regulation of  $\text{Ca}^{2+}_e$ . Unlike many GPCRs,  
 231 the CaSR is chronically exposed to its ligand at baseline concentrations, and thus the receptor  
 232 undergoes very little functional desensitisation (Gama and Breitwieser 1998). Furthermore, CaSR is  
 233 able to elicit signalling responses for as long as elevated  $\text{Ca}^{2+}_e$  is available (Grant, et al. 2011; Lorenz  
 234 et al. 2007; Pi et al. 2005). Experimental studies have sought to explain this apparent paradox and  
 235 using a combination of total internal reflection fluorescence microscopy (TIRFm) and a construct that  
 236 allows the simultaneous measurement of CaSR insertion within plasma membranes and endocytosis  
 237 (known as BSEP-CaSR), a new model for CaSR cell surface expression has emerged (Fig. 4). This  
 238 model proposes that in the basal state, CaSR is only weakly expressed at the plasma membrane, but  
 239 on exposure to increases in  $\text{Ca}^{2+}_e$  there is an increase in anterograde trafficking through the secretory  
 240 pathway (Grant et al. 2011) (Fig. 4). This increase in the secretory pathway involves both mature  
 241 CaSR located in pre-plasma membrane regions, and newly synthesised CaSR from the ER  
 242 (Breitwieser 2013; Grant et al. 2011), and at present the mechanisms that trigger the ADIS events are  
 243 not fully elucidated. However, studies in which ADIS and CaSR signalling were measured  
 244 simultaneously revealed that elevated  $\text{Ca}^{2+}_i$  and the  $\text{G}\alpha_{q/11}$  pathway is likely involved (Gorvin et al.  
 245 2018b; Grant, et al. 2012). Furthermore, the 14-3-3 proteins, which bind CaSR at an Arg-rich site  
 246 within the C-terminus, limits the release of CaSR from the endoplasmic reticulum, and therefore  
 247 regulates CaSR cell surface expression by reducing the ADIS-mobilisable pool of the receptor (Grant,  
 248 et al. 2015). The presence of the ADIS system and constitutive endocytosis may also explain some of  
 249 the inconsistencies between studies of CaSR trafficking, as endocytosis of the receptor is balanced by  
 250 the continual insertion of new CaSR from the secretory pathway (Breitwieser 2013). Those studies in  
 251 which a tagged construct and cell surface labelling (e.g. FLAG) was used in isolation, without also  
 252 measuring insertion of new CaSR within the plasma membrane, may not reflect the physiological  
 253 state. Similarly, measurement of total cell surface expression of receptor cannot be used as a surrogate  
 254 for determining the rate of internalisation.

255

These studies have demonstrated that trafficking of the CaSR plays a critical role in receptor signalling and calcium homeostasis. More recent studies, building upon these initial findings, have identified further insights into CaSR cell surface expression and endocytosis, and recognised further connections between the trafficking and signalling of this receptor.

### 3.1.3 Regulation of cell surface expression by endocytosis

The CaSR has been described to have two types of internalisation: a constitutive pathway (Grant et al. 2011), and an agonist-driven pathway (Gorvin et al. 2018b; Lorenz et al. 2007). The CaSR was shown to internalise using clathrin-mediated endocytosis in early studies (Holstein, et al. 2004). In addition, CaSR has been described to associate with several proteins that facilitate clustering at the plasma membrane and therefore increase the efficiency of internalisation, or act as scaffolds to enable signalling to occur. One such protein is caveolin-1, with which CaSR has been shown to coimmunoprecipitate (Kifor, et al. 1998). CaSR has been described to be enriched in caveolae structures in parathyroid chief cells and osteosarcoma cells (Jung, et al. 2005; Kifor et al. 1998; Sun and Murphy 2010). CaSR-mediated signalling is impaired in cells treated with caveolin-1 targeting siRNA, and it is likely that this signalling function is facilitated by caveolae acting as signalling hubs allowing the CaSR, G-proteins and PKC to cluster together (Jung et al. 2005; Kifor et al. 1998; Sun and Murphy 2010). However, it is currently unknown whether caveolin facilitates CaSR clustering within caveolae structures to enhance signalling, or whether the signalling itself drives this clustering to encourage endocytosis (Breitwieser 2013). Filamin is an actin binding cytoskeleton protein that is important for protein scaffolding (Hjalm, et al. 2001) and may also facilitate clustering that aids in receptor endocytosis. Filamin binds to the CaSR C-terminus and increases total cellular content of CaSR by preventing its proteosomal degradation (Hjalm et al. 2001), with some studies showing CaSR expression is reduced in cells treated with siRNA targeted against filamin. However, other reports show no such changes in CaSR expression, and this likely requires further investigation (Huang, et al. 2006; Mingione, et al. 2017). The net result of Filamin A binding to CaSR is increased MAPK signalling by the receptor (Hjalm et al. 2001; Pi, et al. 2002).

283

284 **3.1.4 Effect of AP2 $\sigma$  mutations on CaSR signalling and trafficking**

285 Mutations in AP2 $\sigma$  have been demonstrated to cause FHH3, and are associated with impaired CaSR-  
 286 mediated Ca<sup>2+</sup><sub>i</sub> signalling (Nesbit et al. 2013b). The AP2 complex is a ubiquitously-expressed  
 287 heterotetrameric protein which plays a fundamental role in the clathrin-mediated endocytosis of  
 288 transmembrane proteins, such as GPCRs. The two larger subunits,  $\alpha$  and  $\beta$ , have appendages that bind  
 289 to the clathrin coat proteins, plasma membrane phospholipids, and endocytic accessory proteins (e.g.  
 290  $\beta$ -arrestin); while the two smaller subunits,  $\mu$  and  $\sigma$ , bind to endocytic motifs of cargo proteins  
 291 (Collins, et al. 2002; Jackson, et al. 2010; Kelly, et al. 2008; Kirchhausen, et al. 2014) (Fig. 4). The  
 292 AP2 $\mu$  subunit recognises tyrosine-based motifs and the AP2 $\sigma$  subunit recognises dileucine-based  
 293 motifs (Haucke and De Camilli 1999; Kelly et al. 2008) (Fig. 4). FHH3-associated mutations in the  
 294 AP2 $\sigma$  protein have been reported to affect the Arg15 residue, and structural modelling studies using a  
 295 published structure of the AP2 complex, have shown these missense mutations (to Cys15, His15 and  
 296 Leu15) likely disrupt interactions with a putative dileucine motif in the CaSR C-terminus (Nesbit et  
 297 al. 2013b), and thus affect CaSR endocytosis.

298

299 Initially this hypothesis was tested using an ELISA assay of total CaSR at cell surfaces in HEK293  
 300 cells stably overexpressing CaSR (HEK-CaSR), and transiently transfected with AP2 $\sigma$ -mutant  
 301 proteins. This showed that AP2 $\sigma$ -mutant expressing cells had increased CaSR cell surface expression  
 302 when compared to AP2 $\sigma$ -WT cells following stimulation with 5mM Ca<sup>2+</sup><sub>e</sub> (Nesbit et al. 2013b). Thus,  
 303 it was concluded that CaSR endocytosis is impaired in these cells resulting in increased total CaSR  
 304 cell surface expression (Nesbit et al. 2013b). More detailed studies characterising the ADIS and  
 305 endocytosis components of CaSR regulation in AP2 $\sigma$  mutant cells using TIRFm and the BSEP-CaSR  
 306 construct, showed both ADIS and CaSR endocytosis were impaired resulting in the net effect of an  
 307 increased total CaSR cell surface expression in cells expressing AP2 $\sigma$  mutant protein compared to  
 308 WT cells (Gorvin et al. 2018b). Additional TIRFm studies focussing on CaSR and clathrin

demonstrated that CaSR and clathrin colocalise at plasma membranes, and that the duration of this colocalisation is prolonged in AP2 $\sigma$ -mutant expressing cells (Gorvin et al. 2018b). Furthermore, the vesicles containing both CaSR and clathrin in AP2 $\sigma$ -mutant cells, when compared to WT cells, were less motile, which is an indication that the vesicles are less likely to result in viable endocytic events (Gorvin et al. 2018b; Rappoport and Simon 2003). Thus, cells expressing AP2 $\sigma$ -mutant proteins have delayed recruitment of clathrin, and colocalisation with CaSR is prolonged, resulting in impaired CaSR endocytosis (Gorvin et al. 2018b).

The effect of the FHH3-associated AP2 $\sigma$ -mutant proteins on multiple CaSR-mediated signalling pathways has been characterised in a number of cellular assays in HEK293 cells stable overexpressing AP2 $\sigma$  WT and mutant proteins and lymphoblastoid cell-lines derived from blood samples from FHH3 patients with the AP2 $\sigma$ -Cys15 mutation (Gorvin et al. 2018b; Nesbit et al. 2013b). These studies confirmed that AP2 $\sigma$  mutations of the Arg15 residue impair  $\text{Ca}^{2+}_i$  mobilisation, phosphorylated ERK1/2 (pERK1/2) MAPK signalling, membrane ruffling and suppression of cAMP, and that all of these pathways occur downstream of  $\text{G}\alpha_{q/11}$  and  $\text{G}\alpha_{i/o}$  (Gorvin et al. 2018b; Nesbit et al. 2013b). Therefore, these studies have shown that AP2 $\sigma$  mutations reduce endocytosis resulting in increased CaSR cell surface expression, but paradoxically decrease CaSR-mediated signalling (Gorvin et al. 2018b). To explain this paradox, we hypothesised that CaSR may be able to continue signalling from within the cell (i.e. sustained signalling) (Gorvin et al. 2018b). Such sustained signalling has been previously reported for some class A (e.g.  $\beta$ 2-adrenergic receptor ( $\beta$ 2AR), dopamine receptor D1 (DRD1), thyroid-stimulating hormone receptor (TSHR), vasopressin receptor 2 (V2R) and luteinizing hormone receptor (LHR)), and class B (e.g. parathyroid hormone 1 receptor (PTH1R)) GPCRs (Calebiro, et al. 2009; Feinstein, et al. 2013; Ferrandon, et al. 2009; Irannejad, et al. 2013; Jean-Alphonse, et al. 2014; Kotowski, et al. 2011). Thus, in cells with the AP2 $\sigma$  mutation, in which there is impaired endocytosis, the availability of internalised receptors from which sustained signals could emanate would be reduced and thus the net effect would be impaired overall CaSR-mediated signalling. To test this hypothesis, a combination of imaging and biochemical analyses, along with

chemical inhibitors were used in HEK-CaSR, HEK-AP2 $\sigma$  and CRISPR-Cas generated  $\beta$ -arrestin knockout cells to assess CaSR-mediated MAPK signalling (Gorvin et al. 2018b). To assess sustained signalling two primary assays were used: 1) assessment of pERK1/2 over 60 minutes following treatment of cells with a 5 min pulse of 5mM Ca<sup>2+</sup><sub>e</sub>; and 2) analysis of SRE luciferase reporter responses over 12 hours, following a 5 min pulse of 5mM Ca<sup>2+</sup><sub>e</sub>. Using these methods, MAPK sustained signals were demonstrated in HEK-CaSR cells, and evidence for an internal, likely endosomal, source was shown in three ways (Gorvin et al. 2018b) (Fig. 5). First, addition of Dyngo, a chemical inhibitor of dynamin, which is required for vesicle scission during clathrin-mediated endocytosis, abolishes sustained signals, whilst rapid plasma membrane mediated signals remain intact. Loss of this sustained response in Dyngo-treated cells was not due to increased apoptosis or decreased proliferation, and was unaffected by inhibition of CaSR protein synthesis as the sustained rise in pERK1/2 was not blocked by tunicamycin, an inhibitor of glycosylation of newly synthesised CaSR. Second, sustained signals were reduced in cells expressing a dominant-negative Rab5 protein, which delays maturation of early endosomes and therefore slows clathrin-mediated endocytosis. Third, reduction or loss of these sustained signals was observed in cells expressing AP2 $\sigma$  mutant proteins (Fig. 5).

The G-proteins involved in these sustained signals were also explored as MAPK signalling is a convergence pathway for multiple CaSR-mediated signalling pathways (Fig. 2). The ability of the G $\alpha_{q/11}$  and G $\alpha_{i/o}$  pathways to activate CaSR sustained signals was investigated using G-protein specific inhibitors (UBO-QIC and PTx) and the SRE luciferase reporter assay, which showed that inhibition of both G $\alpha_{q/11}$  and G $\alpha_{i/o}$  impaired the early, plasma membrane mediated CaSR signal, while only G $\alpha_{q/11}$  was important for the later sustained signal. Furthermore, confocal microscopy confirmed that G $\alpha_q$  and phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>), the lipid hydrolysed by PLC, colocalise with CaSR at plasma membranes and a subpopulation of endosomes (Gorvin et al. 2018b). In addition, inhibitors of the PLC-DAG-IP<sub>3</sub> pathway (U73122, GF-109203X and 2-aminoethoxydiphenyl borate (2-APB), which inhibit PLC, PKC and the IP<sub>3</sub>-receptor, respectively)

were shown to reduce sustained pERK1/2 signals, indicating that  $G\alpha_{q/11}$  and its signalling pathway are important for CaSR-mediated signalling (Gorvin et al. 2018b). Finally, the effects of the  $\beta$ -arrestin scaffold proteins, which are important for the sustained endosomal signalling of some GPCRs such as V2R and PTH1R (Feinstein et al. 2013; Wehbi, et al. 2013), were assessed in HEK-CaSR cells with deletion of the  $\beta$ -arrestin1 and  $\beta$ -arrestin2 genes. In these cells both pERK and SRE reporter responses were unaffected by deletion of the  $\beta$ -arrestin proteins (Gorvin et al. 2018b). Thus, the CaSR mediates some MAPK signals from endosomes using  $G\alpha_q$  and PLC, but does not require  $\beta$ -arrestin for this pathway.

### **Conclusions and important lessons for the future (or unanswered questions)**

Since the cloning of the CaSR gene 25 years ago (Brown, et al. 1993), many insights have been gained into the role of the CaSR in calcium homeostasis, and by studying patients with mutations within this gene we have learnt much about CaSR signalling and trafficking mechanisms. The discovery that CaSR can signal from within the cell, most likely from an endosomal source, and that disruption of a specific structural motif can mediate G-protein independent signalling bias opens up many new avenues of investigation. How do mutations in the CaSR affect these endosomal pathways? Can drugs that target the CaSR affect these pathways? How are trafficking and signalling of the CaSR so intimately linked? Future studies investigating these questions will undoubtedly reveal further complexities into the regulation of CaSR, and could provide mechanisms relevant to other GPCRs.

**Declaration of interest:** The author has no conflict of interest.

**Funding:** This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.



389 **References**

- 390 Ballesteros JA, Jensen AD, Liapakis G, Rasmussen SG, Shi L, Gether U & Javitch JA 2001  
 391 Activation of the beta 2-adrenergic receptor involves disruption of an ionic lock between the  
 392 cytoplasmic ends of transmembrane segments 3 and 6. *J Biol Chem* **276** 29171-29177.
- 393 Bouschet T, Martin S & Henley JM 2005 Receptor-activity-modifying proteins are required for  
 394 forward trafficking of the calcium-sensing receptor to the plasma membrane. *J Cell Sci* **118** 4709-  
 395 4720.
- 396 Breitwieser GE 2012 Minireview: the intimate link between calcium sensing receptor trafficking and  
 397 signaling: implications for disorders of calcium homeostasis. *Mol Endocrinol* **26** 1482-1495.
- 398 Breitwieser GE 2013 The calcium sensing receptor life cycle: trafficking, cell surface expression, and  
 399 degradation. *Best Pract Res Clin Endocrinol Metab* **27** 303-313.
- 400 Brown EM 1991 Extracellular Ca<sup>2+</sup> sensing, regulation of parathyroid cell function, and role of Ca<sup>2+</sup>  
 401 and other ions as extracellular (first) messengers. *Physiol Rev* **71** 371-411.
- 402 Brown EM, Gamba G, Riccardi D, Lombardi M, Butters R, Kifor O, Sun A, Hediger MA, Lytton J &  
 403 Hebert SC 1993 Cloning and characterization of an extracellular Ca(2+)-sensing receptor from bovine  
 404 parathyroid. *Nature* **366** 575-580.
- 405 Calebiro D, Nikolaev VO, Gagliani MC, de Filippis T, Dees C, Tacchetti C, Persani L & Lohse MJ  
 406 2009 Persistent cAMP-signals triggered by internalized G-protein-coupled receptors. *PLoS Biol* **7**  
 407 e1000172.
- 408 Chang W, Tu C, Cheng Z, Rodriguez L, Chen TH, Gassmann M, Bettler B, Margeta M, Jan LY &  
 409 Shoback D 2007 Complex formation with the Type B gamma-aminobutyric acid receptor affects the  
 410 expression and signal transduction of the extracellular calcium-sensing receptor. Studies with HEK-  
 411 293 cells and neurons. *J Biol Chem* **282** 25030-25040.
- 412 Chattopadhyay N & Brown EM 2006 Role of calcium-sensing receptor in mineral ion metabolism and  
 413 inherited disorders of calcium-sensing. *Mol Genet Metab* **89** 189-202.
- 414 Collins BM, McCoy AJ, Kent HM, Evans PR & Owen DJ 2002 Molecular architecture and functional  
 415 model of the endocytic AP2 complex. *Cell* **109** 523-535.
- 416 Conigrave AD & Ward DT 2013 Calcium-sensing receptor (CaSR): pharmacological properties and  
 417 signaling pathways. *Best Pract Res Clin Endocrinol Metab* **27** 315-331.
- 418 Fan G, Goldsmith PK, Collins R, Dunn CK, Krapcho KJ, Rogers KV & Spiegel AM 1997 N-linked  
 419 glycosylation of the human Ca<sup>2+</sup> receptor is essential for its expression at the cell surface.  
 420 *Endocrinology* **138** 1916-1922.
- 421 Feinstein TN, Yui N, Webber MJ, Wehbi VL, Stevenson HP, King JD, Jr., Hallows KR, Brown D,  
 422 Bouley R & Vilardaga JP 2013 Noncanonical control of vasopressin receptor type 2 signaling by  
 423 retromer and arrestin. *J Biol Chem* **288** 27849-27860.
- 424 Ferrandon S, Feinstein TN, Castro M, Wang B, Bouley R, Potts JT, Gardella TJ & Vilardaga JP 2009  
 425 Sustained cyclic AMP production by parathyroid hormone receptor endocytosis. *Nat Chem Biol* **5**  
 426 734-742.
- 427 Firek AF, Kao PC & Heath H, 3rd 1991 Plasma intact parathyroid hormone (PTH) and PTH-related  
 428 peptide in familial benign hypercalcemia: greater responsiveness to endogenous PTH than in primary  
 429 hyperparathyroidism. *J Clin Endocrinol Metab* **72** 541-546.
- 430 Gama L & Breitwieser GE 1998 A carboxyl-terminal domain controls the cooperativity for  
 431 extracellular Ca<sup>2+</sup> activation of the human calcium sensing receptor. A study with receptor-green  
 432 fluorescent protein fusions. *J Biol Chem* **273** 29712-29718.
- 433 Gama L, Wilt SG & Breitwieser GE 2001 Heterodimerization of calcium sensing receptors with  
 434 metabotropic glutamate receptors in neurons. *J Biol Chem* **276** 39053-39059.
- 435 Geng Y, Mosyak L, Kurinov I, Zuo H, Sturchler E, Cheng TC, Subramanyam P, Brown AP, Brennan  
 436 SC, Mun HC, et al. 2016 Structural mechanism of ligand activation in human calcium-sensing  
 437 receptor. *Elife* **5**.
- 438 Gorvin CM, Babinsky VN, Malinauskas T, Nissen PH, Schou AJ, Hanyaloglu AC, Siebold C, Jones  
 439 EY, Hannan FM & Thakker RV 2018a A calcium-sensing receptor mutation causing hypocalcemia  
 440 disrupts a transmembrane salt bridge to activate  $\beta$ -arrestin biased signaling. *Sci Signal* **11**.

- Gorvin CM, Cranston T, Hannan FM, Rust N, Qureshi A, Nesbit MA & Thakker RV 2016 A G-protein Subunit- $\alpha$ 11 Loss-of-Function Mutation, Thr54Met, Causes Familial Hypocalciuric Hypercalcemia Type 2 (FHH2). *J Bone Miner Res* **31** 1200-1206.
- Gorvin CM, Hannan FM, Cranston T, Valta H, Makitie O, Schalin-Jantti C & Thakker RV 2017a Cinacalcet Rectifies Hypercalcemia in a Patient With Familial Hypocalciuric Hypercalcemia Type 2 (FHH2) Caused by a Germline Loss-of-Function  $\alpha$ 11 Mutation. *J Bone Miner Res*.
- Gorvin CM, Hannan FM, Howles SA, Babinsky VN, Piret SE, Rogers A, Freidin AJ, Stewart M, Paudyal A, Hough TA, et al. 2017b  $\alpha$ 11 mutation in mice causes hypocalcemia rectifiable by calcilytic therapy. *JCI Insight* **2** e91103.
- Gorvin CM, Rogers A, Hastoy B, Tarasov AI, Frost M, Sposini S, Inoue A, Whyte MP, Rorsman P, Hanyaloglu AC, et al. 2018b AP2 $\sigma$  mutations impair calcium-sensing receptor trafficking and signaling revealing an endosomal pathway that spatially directs G-protein selectivity. *Cell Reports* **22** 1054-1066.
- Grant MP, Cavanaugh A & Breitwieser GE 2015 14-3-3 Proteins Buffer Intracellular Calcium Sensing Receptors to Constrain Signaling. *PLoS One* **10** e0136702.
- Grant MP, Stepanchick A & Breitwieser GE 2012 Calcium signaling regulates trafficking of familial hypocalciuric hypercalcemia (FHH) mutants of the calcium sensing receptor. *Mol Endocrinol* **26** 2081-2091.
- Grant MP, Stepanchick A, Cavanaugh A & Breitwieser GE 2011 Agonist-driven maturation and plasma membrane insertion of calcium-sensing receptors dynamically control signal amplitude. *Sci Signal* **4** ra78.
- Hannan FM, Howles SA, Rogers A, Cranston T, Gorvin CM, Babinsky VN, Reed AA, Thakker CE, Bockenhauer D, Brown RS, et al. 2015 Adaptor protein-2 sigma subunit mutations causing familial hypocalciuric hypercalcaemia type 3 (FHH3) demonstrate genotype-phenotype correlations, codon bias and dominant-negative effects. *Hum Mol Genet* **24** 5079-5092.
- Hannan FM, Nesbit MA, Zhang C, Cranston T, Curley AJ, Harding B, Fratter C, Rust N, Christie PT, Turner JJ, et al. 2012 Identification of 70 calcium-sensing receptor mutations in hyper- and hypocalcaemic patients: evidence for clustering of extracellular domain mutations at calcium-binding sites. *Hum Mol Genet* **21** 2768-2778.
- Hannan FM & Thakker RV 2013 Calcium-sensing receptor (CaSR) mutations and disorders of calcium, electrolyte and water metabolism. *Best Pract Res Clin Endocrinol Metab* **27** 359-371.
- Haucke V & De Camilli P 1999 AP-2 recruitment to synaptotagmin stimulated by tyrosine-based endocytic motifs. *Science* **285** 1268-1271.
- Hjalm G, MacLeod RJ, Kifor O, Chattopadhyay N & Brown EM 2001 Filamin-A binds to the carboxyl-terminal tail of the calcium-sensing receptor, an interaction that participates in CaR-mediated activation of mitogen-activated protein kinase. *Journal of Biological Chemistry* **276** 34880-34887.
- Hofer AM & Brown EM 2003 Extracellular calcium sensing and signalling. *Nat Rev Mol Cell Biol* **4** 530-538.
- Holstein DM, Berg KA, Leeb-Lundberg LM, Olson MS & Saunders C 2004 Calcium-sensing receptor-mediated ERK1/2 activation requires  $\alpha$ 2 coupling and dynamin-independent receptor internalization. *J Biol Chem* **279** 10060-10069.
- Hu J, Jiang J, Costanzi S, Thomas C, Yang W, Feyen JH, Jacobson KA & Spiegel AM 2006 A missense mutation in the seven-transmembrane domain of the human  $\text{Ca}^{2+}$  receptor converts a negative allosteric modulator into a positive allosteric modulator. *J Biol Chem* **281** 21558-21565.
- Hu J, McLarnon SJ, Mora S, Jiang J, Thomas C, Jacobson KA & Spiegel AM 2005 A region in the seven-transmembrane domain of the human  $\text{Ca}^{2+}$  receptor critical for response to  $\text{Ca}^{2+}$ . *J Biol Chem* **280** 5113-5120.
- Hu J, Reyes-Cruz G, Chen W, Jacobson KA & Spiegel AM 2002 Identification of acidic residues in the extracellular loops of the seven-transmembrane domain of the human  $\text{Ca}^{2+}$  receptor critical for response to  $\text{Ca}^{2+}$  and a positive allosteric modulator. *J Biol Chem* **277** 46622-46631.
- Huang C, Wu Z, Hujer KM & Miller RT 2006 Silencing of filamin A gene expression inhibits  $\text{Ca}^{2+}$  -sensing receptor signaling. *FEBS Lett* **580** 1795-1800.
- Huang Y, Cavanaugh A & Breitwieser GE 2011 Regulation of stability and trafficking of calcium-sensing receptors by pharmacologic chaperones. *Adv Pharmacol* **62** 143-173.

496 Irannejad R, Tomshine JC, Tomshine JR, Chevalier M, Mahoney JP, Steyaert J, Rasmussen SG,  
 497 Sunahara RK, El-Samad H, Huang B, et al. 2013 Conformational biosensors reveal GPCR signalling  
 498 from endosomes. *Nature* **495** 534-538.  
 499 Jackson LP, Kelly BT, McCoy AJ, Gaffry T, James LC, Collins BM, Honing S, Evans PR & Owen  
 500 DJ 2010 A large-scale conformational change couples membrane recruitment to cargo binding in the  
 501 AP2 clathrin adaptor complex. *Cell* **141** 1220-1229.  
 502 Jean-Alphonse F, Bowersox S, Chen S, Beard G, Puthenveedu MA & Hanyaloglu AC 2014 Spatially  
 503 restricted G protein-coupled receptor activity via divergent endocytic compartments. *J Biol Chem* **289**  
 504 3960-3977.  
 505 Jung SY, Kwak JO, Kim HW, Kim DS, Ryu SD, Ko CB & Cha SH 2005 Calcium sensing receptor  
 506 forms complex with and is up-regulated by caveolin-1 in cultured human osteosarcoma (Saos-2) cells.  
 507 *Experimental and Molecular Medicine* **37** 91-100.  
 508 Kelly BT, McCoy AJ, Spate K, Miller SE, Evans PR, Honing S & Owen DJ 2008 A structural  
 509 explanation for the binding of endocytic dileucine motifs by the AP2 complex. *Nature* **456** 976-979.  
 510 Kifor O, Diaz R, Butters R, Kifor I & Brown EM 1998 The calcium-sensing receptor is localized in  
 511 caveolin-rich plasma membrane domains of bovine parathyroid cells. *J Biol Chem* **273** 21708-21713.  
 512 Kifor O, MacLeod RJ, Diaz R, Bai M, Yamaguchi T, Yao T, Kifor I & Brown EM 2001 Regulation  
 513 of MAP kinase by calcium-sensing receptor in bovine parathyroid and CaR-transfected HEK293 cells.  
 514 *Am J Physiol Renal Physiol* **280** F291-302.  
 515 Kirchhausen T, Owen D & Harrison SC 2014 Molecular structure, function, and dynamics of clathrin-  
 516 mediated membrane traffic. *Cold Spring Harb Perspect Biol* **6** a016725.  
 517 Kotowski SJ, Hopf FW, Seif T, Bonci A & von Zastrow M 2011 Endocytosis promotes rapid  
 518 dopaminergic signaling. *Neuron* **71** 278-290.  
 519 Leach K, Wen A, Cook AE, Sexton PM, Conigrave AD & Christopoulos A 2013 Impact of clinically  
 520 relevant mutations on the pharmacoregulation and signaling bias of the calcium-sensing receptor by  
 521 positive and negative allosteric modulators. *Endocrinology* **154** 1105-1116.  
 522 Leach K, Wen A, Davey AE, Sexton PM, Conigrave AD & Christopoulos A 2012 Identification of  
 523 molecular phenotypes and biased signaling induced by naturally occurring mutations of the human  
 524 calcium-sensing receptor. *Endocrinology* **153** 4304-4316.  
 525 Li D, Opas EE, Tuluc F, Metzger DL, Hou C, Hakonarson H & Levine MA 2014 Autosomal  
 526 dominant hypoparathyroidism caused by germline mutation in GNA11: phenotypic and molecular  
 527 characterization. *J Clin Endocrinol Metab* **99** E1774-1783.  
 528 Lorenz S, Frenzel R, Paschke R, Breitwieser GE & Miedlich SU 2007 Functional desensitization of  
 529 the extracellular calcium-sensing receptor is regulated via distinct mechanisms: role of G protein-  
 530 coupled receptor kinases, protein kinase C and beta-arrestins. *Endocrinology* **148** 2398-2404.  
 531 Mannstadt M, Harris M, Bravenboer B, Chitturi S, Dreijerink KM, Lambright DG, Lim ET, Daly MJ,  
 532 Gabriel S & Juppner H 2013 Germline mutations affecting Galphal1 in hypoparathyroidism. *N Engl J*  
 533 *Med* **368** 2532-2534.  
 534 Marx SJ 2015 Letter to the editor: Distinguishing typical primary hyperparathyroidism from familial  
 535 hypocalciuric hypercalcemia by using an index of urinary calcium. *J Clin Endocrinol Metab* **100** L29-  
 536 30.  
 537 Miedlich SU, Gama L, Seuwen K, Wolf RM & Breitwieser GE 2004 Homology modeling of the  
 538 transmembrane domain of the human calcium sensing receptor and localization of an allosteric  
 539 binding site. *J Biol Chem* **279** 7254-7263.  
 540 Mingione A, Verdelli C, Ferrero S, Vaira V, Guarnieri V, Scillitani A, Vicentini L, Balza G, Beretta  
 541 E, Terranegra A, et al. 2017 Filamin A is reduced and contributes to the CASR sensitivity in human  
 542 parathyroid tumors. *J Mol Endocrinol* **58** 91-103.  
 543 Nesbit MA, Hannan FM, Howles SA, Babinsky VN, Head RA, Cranston T, Rust N, Hobbs MR,  
 544 Heath H, 3rd & Thakker RV 2013a Mutations affecting G-protein subunit alpha11 in hypercalcemia  
 545 and hypocalcemia. *N Engl J Med* **368** 2476-2486.  
 546 Nesbit MA, Hannan FM, Howles SA, Reed AA, Cranston T, Thakker CE, Gregory L, Rimmer AJ,  
 547 Rust N, Graham U, et al. 2013b Mutations in AP2S1 cause familial hypocalciuric hypercalcemia type  
 548 3. *Nat Genet* **45** 93-97.

- 549 Obermannova B, Banghova K, Sumnik Z, Dvorakova HM, Betka J, Fencel F, Kolouskova S, Cinek O  
 550 & Lebl J 2009 Unusually severe phenotype of neonatal primary hyperparathyroidism due to a  
 551 heterozygous inactivating mutation in the CASR gene. *Eur J Pediatr* **168** 569-573.
- 552 Pearce SH, Bai M, Quinn SJ, Kifor O, Brown EM & Thakker RV 1996a Functional characterization  
 553 of calcium-sensing receptor mutations expressed in human embryonic kidney cells. *J Clin Invest* **98**  
 554 1860-1866.
- 555 Pearce SH, Trump D, Wooding C, Besser GM, Chew SL, Grant DB, Heath DA, Hughes IA, Paterson  
 556 CR, Whyte MP, et al. 1995 Calcium-sensing receptor mutations in familial benign hypercalcemia and  
 557 neonatal hyperparathyroidism. *J Clin Invest* **96** 2683-2692.
- 558 Pearce SH, Williamson C, Kifor O, Bai M, Coulthard MG, Davies M, Lewis-Barned N, McCredie D,  
 559 Powell H, Kendall-Taylor P, et al. 1996b A familial syndrome of hypocalcemia with hypercalciuria  
 560 due to mutations in the calcium-sensing receptor. *N Engl J Med* **335** 1115-1122.
- 561 Petrel C, Kessler A, Dauban P, Dodd RH, Rognan D & Ruat M 2004 Positive and negative allosteric  
 562 modulators of the Ca<sup>2+</sup>-sensing receptor interact within overlapping but not identical binding sites in  
 563 the transmembrane domain. *J Biol Chem* **279** 18990-18997.
- 564 Pi M, Oakley RH, Gesty-Palmer D, Cruickshank RD, Spurney RF, Luttrell LM & Quarles LD 2005  
 565 Beta-arrestin- and G protein receptor kinase-mediated calcium-sensing receptor desensitization. *Mol*  
 566 *Endocrinol* **19** 1078-1087.
- 567 Pi M, Spurney RF, Tu QS, Hinson T & Quarles LD 2002 Calcium-sensing receptor activation of Rho  
 568 involves filamin and Rho-guanine nucleotide exchange factor. *Endocrinology* **143** 3830-3838.
- 569 Pidasheva S, Grant M, Canaff L, Ercan O, Kumar U & Hendy GN 2006 Calcium-sensing receptor  
 570 dimerizes in the endoplasmic reticulum: biochemical and biophysical characterization of CASR  
 571 mutants retained intracellularly. *Hum Mol Genet* **15** 2200-2209.
- 572 Piret SE, Gorvin CM, Pagnamenta AT, Howles SA, Cranston T, Rust N, Nesbit MA, Glaser B, Taylor  
 573 JC, Buchs AE, et al. 2016 Identification of a G-Protein Subunit- $\alpha$ 11 Gain-of-Function Mutation,  
 574 Val340Met, in a Family With Autosomal Dominant Hypocalcemia Type 2 (ADH2). *J Bone Miner*  
 575 *Res* **31** 1207-1214.
- 576 Pollak MR, Brown EM, Chou YH, Hebert SC, Marx SJ, Steinmann B, Levi T, Seidman CE &  
 577 Seidman JG 1993 Mutations in the human Ca(2+)-sensing receptor gene cause familial hypocalciuric  
 578 hypercalcemia and neonatal severe hyperparathyroidism. *Cell* **75** 1297-1303.
- 579 Rappoport JZ & Simon SM 2003 Real-time analysis of clathrin-mediated endocytosis during cell  
 580 migration. *J Cell Sci* **116** 847-855.
- 581 Rasmussen SG, Choi HJ, Rosenbaum DM, Kobilka TS, Thian FS, Edwards PC, Burghammer M,  
 582 Ratnala VR, Sanishvili R, Fischetti RF, et al. 2007 Crystal structure of the human beta2 adrenergic G-  
 583 protein-coupled receptor. *Nature* **450** 383-387.
- 584 Riccardi D & Brown EM 2010 Physiology and pathophysiology of the calcium-sensing receptor in  
 585 the kidney. *Am J Physiol Renal Physiol* **298** F485-499.
- 586 Schrage R, Schmitz AL, Gaffal E, Annala S, Kehraus S, Wenzel D, Bullesbach KM, Bald T, Inoue A,  
 587 Shinjo Y, et al. 2015 The experimental power of FR900359 to study Gq-regulated biological  
 588 processes. *Nat Commun* **6** 10156.
- 589 Shukla AK, Westfield GH, Xiao K, Reis RI, Huang LY, Tripathi-Shukla P, Qian J, Li S, Blanc A,  
 590 Oleskie AN, et al. 2014 Visualization of arrestin recruitment by a G-protein-coupled receptor. *Nature*  
 591 **512** 218-222.
- 592 Shukla AK, Xiao K & Lefkowitz RJ 2011 Emerging paradigms of beta-arrestin-dependent seven  
 593 transmembrane receptor signaling. *Trends Biochem Sci* **36** 457-469.
- 594 Standfuss J, Edwards PC, D'Antona A, Fransen M, Xie G, Oprian DD & Schertler GF 2011 The  
 595 structural basis of agonist-induced activation in constitutively active rhodopsin. *Nature* **471** 656-660.
- 596 Sun J & Murphy E 2010 Calcium-sensing receptor: a sensor and mediator of ischemic  
 597 preconditioning in the heart. *Am J Physiol Heart Circ Physiol* **299** H1309-1317.
- 598 Takasaki J, Saito T, Taniguchi M, Kawasaki T, Moritani Y, Hayashi K & Kobori M 2004 A novel  
 599 Galphaq/11-selective inhibitor. *J Biol Chem* **279** 47438-47445.
- 600 Thomsen AR, Hvidtfeldt M & Brauner-Osborne H 2012 Biased agonism of the calcium-sensing  
 601 receptor. *Cell Calcium* **51** 107-116.

Uckun-Kitapci A, Underwood LE, Zhang J & Moats-Staats B 2005 A novel mutation (E767K) in the second extracellular loop of the calcium sensing receptor in a family with autosomal dominant hypocalcemia. *Am J Med Genet A* **132A** 125-129.

Vargas-Poussou R, Huang C, Hulin P, Houillier P, Jeunemaitre X, Paillard M, Planelles G, Dechaux M, Miller RT & Antignac C 2002 Functional characterization of a calcium-sensing receptor mutation in severe autosomal dominant hypocalcemia with a Bartter-like syndrome. *J Am Soc Nephrol* **13** 2259-2266.

Volpe A, Guerriero A, Marchetta A, Caramaschi P & Furlani L 2009 Familial hypocalciuric hypercalcemia revealed by chondrocalcinosis. *Joint Bone Spine* **76** 708-710.

Ward DT, Brown EM & Harris HW 1998 Disulfide bonds in the extracellular calcium-polyvalent cation-sensing receptor correlate with dimer formation and its response to divalent cations in vitro. *J Biol Chem* **273** 14476-14483.

Watanabe S, Fukumoto S, Chang H, Takeuchi Y, Hasegawa Y, Okazaki R, Chikatsu N & Fujita T 2002 Association between activating mutations of calcium-sensing receptor and Bartter's syndrome. *Lancet* **360** 692-694.

Wehbi VL, Stevenson HP, Feinstein TN, Calero G, Romero G & Vilardaga JP 2013 Noncanonical GPCR signaling arising from a PTH receptor-arrestin-Gbetagamma complex. *Proc Natl Acad Sci U S A* **110** 1530-1535.

Wu H, Wang C, Gregory KJ, Han GW, Cho HP, Xia Y, Niswender CM, Katritch V, Meiler J, Cherezov V, et al. 2014 Structure of a class C GPCR metabotropic glutamate receptor 1 bound to an allosteric modulator. *Science* **344** 58-64.

Zhang C, Zhang T, Zou J, Miller CL, Gorkhali R, Yang JY, Schilmiller A, Wang S, Huang K, Brown EM, et al. 2016 Structural basis for regulation of human calcium-sensing receptor by magnesium ions and an unexpected tryptophan derivative co-agonist. *Sci Adv* **2** e1600241.

1     **Figures**

2     **Figure 1           Regulation of extracellular calcium at the parathyroid glands**

3     **(A)** Schematic illustrating how the parathyroid glands respond to changes in extracellular calcium  
4     concentrations  $[Ca^{2+}_e]$  in the blood. In the presence of low  $[Ca^{2+}_e]$  in the blood, the parathyroid glands  
5     secrete parathyroid hormone (PTH). PTH binds to PTH receptors at the kidney, resulting in calcium  
6     reabsorption, and activation of 1,25-dihydroxyvitamin D<sub>3</sub>; which in turn acts upon bone to mediate  
7     calcium release, and the intestine to activate calcium resorption. PTH also acts directly on bone to  
8     activate calcium release. The tissue-specific effects on calcium are shown in the hatched box with an  
9     arrow to indicate that these are increased by low  $[Ca^{2+}]$  in the blood. **(B)** Schematic of the  
10    homodimeric calcium-sensing receptor (CaSR) showing the two protomers (CaSR1 and CaSR2)  
11    within the plasma membrane (PM). The calcium-sensing receptor is expressed highly at the  
12    parathyroid gland and its major function is to detect  $[Ca^{2+}_e]$  and regulate PTH secretion accordingly.  
13    The CaSR has a large extracellular domain (ECD) comprising the venus flytrap domain (VFTD) that  
14    contains two lobes (lobe 1 and lobe 2), and a cysteine-rich domain (CRD). The CRD connects the  
15    ECD to the 7 transmembrane domains (TMD), and the CaSR has a long cytoplasmic intracellular  
16    domain (ICD). Ionised calcium binds between the two lobes of the VFTD. In the presence of high  
17    calcium the CaSR is activated, leading to changes in its signal transduction and reduced PTH  
18    secretion.

19    **Figure 2           Major signalling pathways of the calcium-sensing receptor (CaSR)**

20    Schematic diagram of the calcium-bound homodimeric CaSR and its major signalling pathways. The  
21    CaSR is expressed at the plasma membrane (PM) where its function is to detect  $[Ca^{2+}_e]$  in the blood.  
22    The CaSR activates two major signalling pathways:  $G_{q/11}$  and  $G_{i/o}$ . Activation of the  $G_{q/11}$  pathway  
23    leads to stimulation of its major effector protein phospholipase C (PLC), which hydrolyses  
24    phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>) to the second messengers diacylglycerol (DAG) and  
25    inositol trisphosphate (IP<sub>3</sub>). DAG activates protein kinase C (PKC), and the mitogen-activated protein  
26    kinase (MAPK) signalling pathway. IP<sub>3</sub> binds to IP<sub>3</sub> receptors at the endoplasmic reticulum (ER),

which mobilise intracellular calcium ( $\text{Ca}^{2+}_i$ ) release into the cytoplasm. This increase in  $\text{Ca}^{2+}_i$  can further activate MAPK signalling. The CaSR also activates  $G_{i/o}$  signalling pathways, which inhibits adenylate cyclase (AC), resulting in reductions in cAMP and protein kinase A (PKA) activity. This reduction in PKA relieves the inhibition on MAPK signalling and therefore provides another activation pathway for MAPK signalling. The net effect of all these signalling pathways is a change in gene transcription.

**Figure 3      Disruption of an Arg680-Glu767 salt-bridge within CaSR mediates signalling bias by activation of a G-protein independent,  $\beta$ -arrestin pathway**

(A) Schematic diagram of a CaSR monomer at the plasma membrane (PM) showing the seven transmembrane domains (TM1-7) with extracellular loops 1–3 (ECL1–3) and intracellular loops 1–3 (ICL1–3). Arg680 is located at the extracellular end of TM3 and is predicted to form a salt-bridge with either a Glu767 residue on ECL2 or a Glu837 residue on TM7. (B) Homology model of the CaSR TM3, TM7 and ECL2 region, reproduced with permission from Gorvin *et al*, 2018, Science Signaling (Gorvin et al. 2018a). The homology model is based on the published structure of mGluR1 (Wu, et al. 2014). The Arg680 residue is shown projecting from TM3 and is predicted to form a salt bridge with Glu767 on ECL2. The Glu837 residue on TM7 lies at a 5.7Å distance from the Arg680 residue, and therefore formation of a salt bridge between Arg680 and Glu837 is less likely, but was tested as the homology model may not reflect the true state of CaSR.

**Figure 4      Mechanisms by which calcium-sensing receptor (CaSR) expression is regulated at plasma membranes**

Schematic showing trafficking pathways regulating the expression of CaSR at the plasma membrane. The CaSR is continuously synthesised at the endoplasmic reticulum (ER) and undergoes post-translational modifications at the ER and Golgi, before export to the plasma membrane surface. In addition to this constitutive anterograde trafficking of the CaSR, an additional export pathway for the CaSR has been described. This pathway is activated by high  $[\text{Ca}^{2+}]_e$  and has been named the agonist-

53 driven insertional signalling, or ADIS, pathway. Removal of the CaSR from the cell surface is  
54 regulated by clathrin-mediated endocytosis that requires the heterotetrameric adaptor protein-2 (AP2).  
55 AP2 binds directly to transmembrane proteins using its  $\mu$ -subunit or  $\sigma$ -subunit. The CaSR has a  
56 putative dileucine motif within its C-terminus with which it is predicted to bind to AP2 $\sigma$ . The  $\alpha$ -  
57 subunit and  $\beta$ -subunit have large appendages that are important for binding to clathrin coat proteins,  
58 plasma membrane phospholipids, and endocytic accessory proteins. Once internalised within clathrin-  
59 coated vesicles, CaSR is targeted to the endosomal-lysosomal system. Experimental evidence  
60 suggests CaSR is degraded at the lysosome, and very little, if any, recycling of the receptor occurs.

61 **Figure 5      The calcium-sensing receptor (CaSR) can activate plasma membrane and**  
62 **endosomal signalling pathways**

63 Schematic diagram showing the signalling pathways that occur downstream of the CaSR. As  
64 described in Figure 2, CaSR predominantly signals via: the  $G_{q/11}$  pathway, leading to activation of the  
65 intracellular calcium ( $Ca^{2+}_i$ ) and mitogen-activated protein kinase (MAPK) pathways; and the  $G_{i/o}$   
66 pathway leading to activation of MAPK and reductions in the cAMP pathways. These pathways are  
67 activated rapidly (within 2-5 minutes), and originate at the plasma membrane. The CaSR can also  
68 signal from an internal location using a  $G_{q/11}$  pathway. This sustained signal that occurs later than the  
69 plasma membrane signal (from 30 minutes), is sensitive to: global clathrin-mediated endocytosis  
70 blockade (shown experimentally using dyngo); inhibition of CaSR-mediated internalisation  
71 (demonstrated in cells expressing FHH3-associated AP2 $\sigma$  mutants); and maturation of internalised  
72 vesicles to the early endosome (demonstrated experimentally using a dominant-negative (DN) form of  
73 Rab5). Thus, this sustained signal is likely to arise within endosomes.

74








Figure 1

78x27mm (300 x 300 DPI)



94x73mm (300 x 300 DPI)

**A**



**B**






Figure 3

109x173mm (300 x 300 DPI)



104x68mm (300 x 300 DPI)



87x42mm (300 x 300 DPI)